

LongRunMIP - motivation and design for a large collection of millennial-length AO-GCM simulations

Article

Accepted Version

Rugenstein, M., Bloch-Johnson, J., Abe-Ouchi, A., Andrews, T., Beyerle, U., Cao, L., Chadha, T., Danabasoglu, G., Dufresne, J.-L., Duan, L., Foujols, M.-A., Frolicher, T., Geoffroy, O., Gregory, J., Knutti, R., Li, C., Marzocchi, A., Mauritsen, T., Menary, M., Moyer, E., Nazarenko, L., Paynter, D., Saint-Martin, D., Schmidt, G. A., Yamamoto, A. and Yang, S. (2020) LongRunMIP - motivation and design for a large collection of millennial-length AO-GCM simulations. *Bulletin of the American Meteorological Society*, 100 (12). pp. 2551-2569. ISSN 1520-0477 doi: <https://doi.org/10.1175/BAMS-D-19-0068.1> Available at <https://centaur.reading.ac.uk/85458/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1175/BAMS-D-19-0068.1>

Publisher: American Meteorological Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in

the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **LongRunMIP – motivation and design for a large collection of**
2 **millennial-length AO-GCM simulations**

3 Maria Rugenstein*

4 *Institute for Atmospheric and Climate Science, ETH Zurich, CH-8092 Zurich, Switzerland*

5 *Max-Planck-Institute for Meteorology, Bundesstrasse 53, 20146 Hamburg, Germany*

6 Jonah Bloch-Johnson

7 *NCAS, University of Reading, Reading*

8 Ayako Abe-Ouchi

9 *Atmosphere and Ocean Research Institute, The University of Tokyo*

10 Timothy Andrews

11 *Met Office Hadley Centre, FitzRoy Road, Exeter, EX1 3PB*

12 Urs Beyerle

13 *Institute for Atmospheric and Climate Science, ETH Zurich, CH-8092 Zurich, Switzerland*

14 Long Cao

15 *School of Earth Sciences, Zhejiang University, Hang Zhou, Zhejiang Province, 310027, China*

16 Tarun Chadha

17 *ITS Research Informatics, ETH Zurich, Switzerland*

18 Gokhan Danabasoglu

19 *National Center for Atmospheric Research, P.O. Box 3000, Boulder, CO 80307*

20 Jean-Louis Dufresne

21 *Centre National de la Recherche Scientifique, Universit Pierre et Marie Curie, ENS, Ecole*

22 *Polytechnique, Paris, France*

23 Lei Duan

24 *School of Earth Sciences, Zhejiang University, Hang Zhou, Zhejiang Province, 310027, China*

25 Marie-Alice Foujols

26 *Institut Pierre-Simon-Laplace, Sorbonne Universit / CNRS, Paris, France*

27 Thomas Frölicher

28 *Climate and Environmental Physics, Physics Institute, University of Bern, Switzerland,*

29 *Oeschger Centre for Climate Change Research, University of Bern, Switzerland*

30 Olivier Geoffroy

31 *CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France*

32 Jonathan Gregory

33 *NCAS, University of Reading, Reading,*

34 *Met Office Hadley Centre, FitzRoy Road, Exeter, EX1 3PB*

35 Reto Knutti

36 *Institute for Atmospheric and Climate Science, ETH Zurich, CH-8092 Zurich, Switzerland*

37 Chao Li

38 *Max-Planck-Institute for Meteorology, Bundesstrasse 53, 20146 Hamburg, Germany*

39 Alice Marzocchi

40 *National Oceanography Centre, European Way, Southampton, SO14 3ZH, UK*

41 Thorsten Mauritsen

42 *Stockholm University, SE-106 91 Stockholm, Sweden*

43 Matthew Menary

44 *LOCEAN, Sorbonne Universit, Paris, France*

45 Elisabeth Moyer

46 *Department of the Geophysical Sciences, University of Chicago, Chicago, Illinois, USA*

47 Larissa Nazarenko

48 *NASA Goddard Institute for Space Studies, 2880 Broadway, New York, NY 10025*

49 David Paynter

50 *Geophysical Fluid Dynamics Laboratory, Princeton, New Jersey, USA*

51 David Saint-Martin

52 *CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France*

53 Gavin A. Schmidt

54 *NASA Goddard Institute for Space Studies, 2880 Broadway, New York, NY 10025*

55 Akitomo Yamamoto

56 *Japan Agency for Marine-Earth Science and Technology, Yokohama, Japan*

57 *Shuting Yang*

58 *Danish Meteorological Institute, Lyngbyvej 100, DK-2100 Copenhagen, Denmark*

59 **Corresponding author address: Maria Rugenstein, Max-Planck-Institute for Meteorology, Bun-*
60 *desstrasse 53, 20146 Hamburg, Germany*

61 *E-mail: maria.rugenstein@mpimet.mpg.de*

ABSTRACT

We present a model intercomparison project, LongRunMIP, the first collection of millennial-length (1000+ year) simulations of complex coupled climate models with a representation of ocean, atmosphere, sea ice, and land surface, and their interactions. Standard model simulations are generally only a few hundred years long. However, modeling the long-term equilibration in response to radiative forcing perturbation is important for understanding many climate phenomena, such as the evolution of ocean circulation, time- and temperature-dependent feedbacks, and the differentiation of forced signal and internal variability. The aim of LongRunMIP is to facilitate research into these questions by serving as an archive for simulations that capture as much of this equilibration as possible. The only requirement to participate in LongRunMIP is to contribute a simulation with elevated, constant CO₂ forcing that lasts at least 1000 years. LongRunMIP is a MIP of opportunity in that the simulations were mostly performed prior to the conception of the archive without an agreed-upon set of experiments. For most models, the archive contains a preindustrial control simulation and simulations with an idealized (typically abrupt) CO₂ forcing. We collect 2D surface and top-of-atmosphere fields, and 3D ocean temperature and salinity fields. Here, we document the collection of simulations and discuss initial results, including the evolution of surface and deep ocean temperature and cloud radiative effects. As of summer 2019, the collection includes 50 simulations of 15 models by 10 modeling centers. The data of LongRunMIP are publicly available. We encourage submission of more simulations in the future.

85 (Capsule Summary) LongRunMIP is the first collection of millennial-length simulations of com-
86 plex coupled climate models and enables investigations of how these models equilibrate in re-
87 sponse to radiative perturbations.

88 1. Motivation and objectives

89 Millennial-length climate simulations are necessary to understand the equilibrium states that oc-
90 cur in response to external forcings, as well as the relationship between transient and equilibrated
91 behavior. Unforced millennial-length simulations are useful as well, as they allow us to consider
92 long-term internal variability and to analyze shorter-term variability with increased statistical cer-
93 tainty. Reasons to study these long time scales include:

- 94 • To better understand long-term climate dynamics. Outstanding issues include the time scales
95 of ocean circulation response (e.g., Jansen et al. 2018; Rind et al. 2018), continental drying
96 trends (e.g., Sniderman et al. 2019), or sea level rise (e.g., Bilbao et al. 2015; Rugenstein et al.
97 2016c).
- 98 • To help predict the impacts of 20th and 21st century emissions on century timescales, such as
99 ice sheet stability, deep ocean warming, or polar amplification (e.g., Frölicher and Joos 2010;
100 Clark et al. 2016; Mauritsen and Pincus 2017), which are rarely explicitly simulated using a
101 fully-coupled climate model.
- 102 • To more accurately estimate Equilibrium Climate Sensitivity (ECS), which is the equilibrium
103 response of the surface air temperature to a doubling of CO₂ due to the “fast” feedbacks water
104 vapor, lapse rate, clouds, and sea ice but excluding Earth system feedbacks such as changes
105 in the carbon cycle, ice sheets, or vegetation. While ECS has long been a focus of scientific

inquiry, substantial uncertainty remains as to its value (e.g., Charney et al. 1979; Knutti et al. 2017).

- To understand the relationship between the transient response of the climate and its equilibration. Since radiative feedbacks can depend on the evolution of the spatial pattern of warming (e.g., Senior and Mitchell 2000; Winton et al. 2010; Armour et al. 2013; Andrews et al. 2015; Andrews and Webb 2018) and on the background temperature (e.g., Colman and McAvaney 2009; Caballero and Huber 2013; Block and Mauritsen 2013; Meraner et al. 2013; Bloch-Johnson et al. 2015), a constant effective sensitivity of the climate is an inadequate assumption. Several methods have been proposed to predict the equilibrium response from transient simulations given a changing global feedback (Held et al. 2010; Winton et al. 2010; Armour et al. 2013; Geoffroy et al. 2013b,a; Frölicher et al. 2014; Proistosescu and Huybers 2017; Saint-Martin et al. 2019), but only fully equilibrated climate model simulations can serve to test how well these methods predict equilibrium conditions.
- To test theories for the relationship between feedbacks at different time-scales (Gregory et al. 2015, 2016; Zhou et al. 2016; Rugenstein et al. 2016a; Armour 2017; Proistosescu and Huybers 2017; Ceppi and Gregory 2017; Andrews and Webb 2018; Andrews et al. 2018), and to quantify the influence of slow, centennial-scale modes on the temperature evolution of the last century (Armour 2017; Proistosescu and Huybers 2017).
- To understand the relevance, time scales, and magnitude of the energy imbalances and drifts exhibited by climate models (e.g., Hobbs et al. 2016), with the potential application of decreasing the spin-up time needed to run these models.
- To understand the relationship between the forced response and internal variability. This relationship is currently studied using the time frame of one or two centuries, which is not

enough to robustly quantify the internal variability under consideration (e.g., Maher et al. 2018; Lutsko and Takahashi 2018; Bloch-Johnson et al. in revision), **millennial time scales with varying forcings** (e.g., Köhler et al. 2017; Khon et al. 2018; Rehfeld et al. 2018) or by using expensive large ensemble simulations **on decadal to centennial time scales** (e.g., Deser et al. 2012; Maher et al. 2019; Rodgers et al. 2015). Millennial long simulations allow us to differentiate the transient response from the equilibrated forced response, even for quantities with large internal variability, such as precipitation, droughts, or the El Niño-Southern Oscillation (ENSO), **and also the significance of a change in internal variability in a transient simulation relative to the control simulation** (e.g., Brown et al. 2017).

- To compare climate model responses and paleo proxies, e.g. of surface or deep ocean temperatures or hydrological conditions on land in order to provide an independent way of testing climate models (Gebbie and Huybers 2019; Burls and Fedorov 2017; Scheff et al. 2017).

With LongRunMIP, we aim to advance knowledge in the above mentioned areas, fill a gap in the CMIP protocols (Taylor et al. 2011; Eyring et al. 2016), and collect published data in one location for easy public access.

The goals of LongRunMIP are

- a) to continuously gather existing millennial-length simulations (both published and unpublished)
- b) to standardize the collected data (e.g., using the same units and sign conventions)
- c) to make the data publicly available and easily accessible
- d) to foster an interdisciplinary community of users working on millennial-length problems, with experts on oceanography, atmospheric dynamics, energy balance modeling, ice sheet modeling, and paleoclimatology

The objectives of this paper are to

- a) motivate the data collection strategy (Section 2)
- b) specify the requirements for LongRunMIP contributors (Section 2 and b)
- c) give an overview of currently submitted simulations and models (Section 2a, b, and Table 1)
- d) give a sample of some initial analysis on these simulations (Section 3)
- e) show how LongRunMIP relates to the existing literature on millennial-length simulations (Section 4a)
- f) discuss the limitations and opportunities of LongRunMIP (Section 4b and c).

2. Experimental design and data collection strategy

LongRunMIP is the first and largest compilation of millennial-length simulations of complex climate models to date, where a “complex climate model” is understood to include an atmospheric, sea ice, land, and full depth ocean component, i.e. Atmosphere-Ocean General Circulation Models (AO-GCMs) with a dynamic atmosphere and ocean, as opposed to Models of Intermediate Complexity (EMICs), which are often used to study millennial-length questions in climate science (e.g., Zickfeld et al. 2013; Levermann et al. 2013). These model simulations include the “fast” feedbacks, such as changes in water vapor, lapse rate, sea ice, and clouds (Charney et al. 1979), but no “slow” feedbacks, such as changes in the ice-sheets. **Vegetation is treated differently in the models (see Section 2b).** In Section 4 we discuss the implications and limitations of **our** approach. Our goal is to collect as many simulations from as many independent models as possible, while keeping the archive and data sharing manageable. Consequently, we keep our requirements for contributions low.

173 *a. Simulations and variables*

174 A step-increase in atmospheric CO₂ concentrations (in the following called “step-forcing”) is
175 one of the simplest experiments for studying a model’s response to forcing and is used as a bench-
176 mark simulation in CMIP3, CMIP5, and CMIP6 (Meehl et al. 2007; Taylor et al. 2011; Eyring
177 et al. 2016). More realistic, gradual forcing scenarios have been shown to be representable by the
178 step-forcing scenarios and exhibit feedbacks that correlate with those computed from step-forcing
179 simulations (Good et al. 2013, 2015; Geoffroy and Saint-Martin 2014; Colman and Hanson 2016).
180 The CMIP3 protocol required a step-forcing of *doubling* atmospheric CO₂ (here referred to as
181 *abrupt2x*) above pre-industrial levels in a slab (i.e. non-dynamical) ocean, which for decades has
182 been used to define ECS (e.g., Charney et al. 1979; Boer and Yu 2003c; Danabasoglu and Gent
183 2009). The integration time scale of these model setups are a couple of decades. However, a
184 *quadrupling* of CO₂ (here referred to as *abrupt4x*) above pre-industrial levels has a better ratio of
185 forced signal to internal variability. Because the forced response was assumed to scale linearly
186 with increased forcing, the CMIP5 protocol requested an abrupt quadrupling of CO₂, now in a
187 fully coupled model with a dynamical ocean, requiring longer integration time scales. The CMIP6
188 protocol again requests abrupt CO₂ quadrupling experiments, but encourages also the submission
189 of abrupt CO₂ doublings, to study the relation between different forcing levels (Eyring et al. 2016;
190 Good et al. 2016). CMIP5 and 6 protocols require the submission of 150 years of model output.
191 A representative response of surface temperature anomalies and top of the atmosphere (TOA) ra-
192 diative imbalance to an *abrupt4x* scenario is shown in Fig. 1. All anomalies mentioned in this
193 paper are computed as the difference of the experiment from the average of the control simulation.
194 After the 150 years of CMIP protocol length (blue shading) and after 1000 years (the minimum
195 contribution to LongRunMIP, light red shading), the surface temperature response of the exem-

196 plary model shown here has reached 75 % and 88 % of its final value respectively, while the TOA
197 radiation has equilibrated 85 % and 93 % of the forcing respectively (7.6 W m^{-2} for this model).
198 Thus, the final equilibration is a CPU-intensive exercise; the model shown here needs 4000 years
199 to balance the final 0.5 W m^{-2} (dark red shading).

200 The set of variables we collect is motivated by the interest of the LongRunMIP contributors and
201 organizers in ECS, temperature and time dependent feedbacks, and deep ocean warming. Table
202 1 lists the variable names, units, and temporal and spatial resolution of the requested variables.
203 The naming and sign conventions follow the CMIP5 protocol¹. Given the large amount of data
204 involved, we have kept our requested variable list low to allow as many groups as possible to
205 participate. For the same reason, we do not request the data to be “CMORized”², i.e. written in
206 conformance with the CMIP standards. However, we do homogenize signs, variable long names,
207 and units, and also provide a regridded version of the fields, as well as global means.

208 *b. Minimal, optimal, and current contributions*

209 The *minimal requirement* to contribute to LongRunMIP are annual fields of a single simulation
210 of any CO_2 forcing scenario that has at least 1000 years of constant forcing, along with a control
211 simulation of any length. The complexity of the model should be CMIP5-class and include dy-
212 namic atmosphere, ocean, and sea ice components. An *optimal contribution* comprises monthly
213 fields of fully equilibrated *abrupt2x*, *4x*, and *8x* simulations and a control simulation of several
214 millennia.

215 Table 2 lists the model characteristics of the *current* contributions. Because the archive is assem-
216 bled from experiments initiated independently for research purposes by multiple modeling groups,
217 there is no pre-defined protocol like for the CMIP simulations. The models are diverse in origin

¹http://cmip-pcmdi.llnl.gov/cmip5/data_description.html

²<https://pcmdi.llnl.gov/CMIP6/Guide/dataUsers.html>

218 and sample the CMIP5 range of models well (see discussion on model genealogy in Knutti 2010).
219 Table 2 lists references for each model and publications using (parts of) the model output. Most
220 of the current contributions to LongRunMIP are extensions of CMIP5 simulations, sometimes
221 with updated model versions, while one model is an extension of a CMIP3 and another model an
222 extension of a CMIP6 contributions (CCSM3 and CNRM-CM6-1 respectively).

223 Many of our current contributions fall short of the optimal expectation for equilibrium, because
224 even several millennia are insufficient for the deep ocean to equilibrate (see discussion around
225 Fig. 4). However, a few millennia appear to be enough for the surface temperature and TOA
226 radiative imbalance to reach a new steady state in most models (see Section 3), and many questions
227 can be adequately addressed with the current contributions. Our approach is to be inclusive, and
228 to leave it to the user to determine the degree of equilibration needed for their research and to
229 develop criteria for model selection.

230 Most contributions are step-forcing simulations, generally to 2x or 4x pre-industrial CO₂ con-
231 centrations (in Fig. 2 abrupt2x in colored in yellow, abrupt4x in orange, abrupt8x in dark red;
232 abrupt2.4x and abrupt4.8x in dark and light pink). There are currently three exceptions: 1) some
233 model simulations have gradual increases in CO₂ at 1% per year until doubled or quadrupled con-
234 centrations are reached, after which the concentration is kept constant (1pct2x and 1pct4x, light
235 and medium red in Fig. 2). 2) One model simulates the 1850-2010 period, after which CO₂ in-
236 creases either piecewise linearly for 90 years until reaching 2.4x pre-industrial values (CCSM3II).
237 3) Finally, one model simulates the historical period and then the CMIP5 extended representative
238 concentration pathway 8.5 (including CH₄, N₂O, CFC11, and CFC12 in addition to CO₂) until
239 year 2300 after which all forcing agents are kept constant (RCP8.5+, violet in Fig. 2) For the
240 models that did not contribute a a millennial-long step-forcing simulation, we collect short (typ-
241 ically 150 year) step-forcing simulations, generally from the CMIP5 archive. These simulations

242 can be used to estimate the effective climate sensitivity and to relate transient and equilibrium
243 responses. They are not mentioned in Table 2 and Fig. 2.

244 Most contributors were able to submit all requested variables. Some models only stored annual
245 output, while for a few models the entire model output (including many more variables than listed
246 in Table 1) is available. In principle, but with considerable effort, additional variables not listed in
247 Table 1 could be requested from some or all contributors.

248 Some models are outliers in some sense. For example, the simulation *abrupt4x* of FAMOUS
249 warms anomalously strong (Fig. 2 and 7) due to a shortwave cloud effect which is positive through-
250 out the simulation and longwave clear-sky effect, which increases anomalously strongly (not
251 shown, see Rugenstein et al. (2019)). In principle though, such extreme behavior could represent
252 possible characteristics of the real world (e.g., Bloch-Johnson et al. 2015; Schneider et al. 2019).
253 Another atypical model is EC-Earth-PISM, which is the only model with an interactive Greenland
254 ice sheet. This additional component and its historical and RCP8.5+ forcing scenario makes it
255 harder to compare the simulation to other models and attribute changes to one forcing component.
256 This model also does not equilibrate but finally produces a negative TOA imbalance, which prob-
257 ably would increase if the simulation was integrated further. We encourage similar “problematic”
258 submissions, since our focus is on understanding model behavior and the large range of model
259 responses (discussed in Section 3).

260 In nine models, the vegetation is fixed to pre-industrial conditions (ECHAM5, CCSM3,
261 CCSM3II, HadCM3L, FAMOUS, MIROC32, ECEARTH, GISSER2R, CNRMCM61), while the
262 other seven models have dynamic vegetation schemes (MPIESM11, MPIESM12, CESM104,
263 HadGEM2, GFDLESM2M, GFDLCM3, IPSLCM5A).

3. Sample of model output

a. Imbalances in the control simulation and drift

In principle, the TOA radiative imbalance should be zero in a control simulation. Most models contributing to LongRunMIP do not loose or gain energy (Fig. 3). However, some models that are equilibrated in the sense that they show no substantial drift, still have a constant energy leakage. For CMIP5 models, imbalances of the same order of magnitude (and larger) have been shown to be uncorrelated with the forced response (Hobbs et al. 2016). If computing atmospheric anomalies, we suggest **users** to take the difference of each time step to the time-averaged control simulation imbalance, except for CCSM3II and GFDL-CM3 for which the difference to a polynomial fit to the control simulation time series seems appropriate (see Fig. 3).

The deep ocean (defined here as **depth level around 2 km**) has an astonishingly small drift in the global average in most models (Fig. 4, lowest panel). While the surface ocean time scales closely follows the global mean surface air temperature anomaly, the deep ocean takes centuries to equilibrate. Panel a and b of Fig. 4 display the surface and deep ocean temperature anomalies, computed as the difference of the forced and control simulations, while the lowest panel shows the absolute temperatures of the deep ocean in the control simulations to indicate the model spread in the base state. Previous work on long-term trends in deep ocean temperature and salinity shows that these trends **may** reflect ongoing changes in stratification and the strength and depth of the Atlantic Meridional Overturning Circulation (AMOC; e.g., Stouffer and Manabe 2003; Rugenstein et al. 2016a; Marzocchi and Jansen 2017; Jansen et al. 2018). Even if the energy flux imbalance at the TOA or the ocean surface are close to a new steady state this does not necessarily indicate that the deep ocean is equilibrated as well (Zhang et al. 2013; Hobbs et al. 2016; Marzocchi and Jansen 2017). Reaching deep ocean equilibration may not be necessary for studies concerned with

287 surface properties only. However, for interpretation of paleo proxies and comparison with model
288 simulations, distinguishing between the transient and equilibrium response in the intermediate or
289 deep ocean is necessary (Zhang et al. 2013; Marzocchi and Jansen 2017; Rind et al. 2018; Jansen
290 et al. 2018).

291 *b. Evolution of surface temperature and cloud radiative effect*

292 The evolution of large scale surface air temperature patterns on decadal to millennial time scales
293 (Fig. 5) are robust among models and different forcing levels. The simulations show a strong land-
294 sea warming contrast on short time scales and little warming over the Southern Ocean on decadal
295 to centennial time scales (e.g., Manabe et al. 1991; Gregory 2000; Joshi and Gregory 2008; Geoff-
296 froy and Saint-Martin 2014; Armour et al. 2016). A warming pattern reminiscent of the positive
297 phase of ENSO and the Interdecadal Pacific Oscillation occurs throughout the Pacific basin (panel
298 b; Held et al. 2010; Song and Zhang 2014; Andrews et al. 2015; Luo et al. 2017) but decays on
299 centennial to millennial time scales (panel c and d), with a large model spread in time scales (not
300 shown). As it approaches equilibrium, the temperature pattern becomes more homogeneous, the
301 land-sea warming contrast reduces (e.g., Held et al. 2010; Geoffroy and Saint-Martin 2014), and
302 the Southern Hemisphere high latitudes keep warming beyond year 1000. As in previous studies,
303 the AMOC first declines (Gregory et al. 2005; Zhu et al. 2014; Kostov et al. 2014; Trossman et al.
304 2016) and then recovers (Stouffer and Manabe 2003; Li et al. 2013; Zickfeld et al. 2013; Rugen-
305 stein et al. 2016a; Rind et al. 2018), resulting in a delayed warming in the North Atlantic. Panel
306 a, b, and e correspond to the blue shading in Fig. 1, and are known from CMIP5 simulations (e.g.,
307 Andrews et al. 2015), while panel c, d, f, and g highlight that the simulations still warm substan-
308 tially on centennial to millennial time scales, mainly in areas with more sensitive – i.e. positive
309 or small negative – feedbacks (Rugenstein et al. 2019). Normalizing the zonal-mean temperature

310 anomaly by the global mean warming reveals the relative **zonal-mean** warming (Fig. 6). Arctic am-
 311 plification begins very early in the simulations and warming throughout the Southern Hemisphere
 312 is lower than the global average in almost all models for the first centuries. Between year 100 and
 313 1000 the Southern Hemisphere warms more than the Northern Hemisphere in all latitudes pole-
 314 ward from 30°, in some regions by more than 4 K. Antarctic warming slowly increases, but is still
 315 substantially less than Arctic amplification (e.g., Salzmann 2017). In a couple of models, the am-
 316 plitude of Antarctic and Arctic amplification is the same after 4000 years of model integration time
 317 (GISSE2R and ECHAM5; Li et al. 2013), while in other models the Antarctic amplification stays
 318 substantially smaller and still increasing after a couple of thousand years. LongRunMIP shows
 319 that there is no reduction in model spread in the polar regions through time and that although all
 320 models follow a similar large scale pattern evolution (Fig. 5), the local response time scales, e.g.
 321 in the North Atlantic, Southern Ocean, or equatorial Pacific differ by hundreds to thousands years.

322 While the large scale temperature response is rather robust between models and simulations,
 323 the cloud radiative effect (CRE) differs strongly in magnitude and time evolution, both between
 324 models and between forcing levels for the same model (Fig 7). We show the shortwave CRE –
 325 computed as the difference between “all sky” and “clear sky” shortwave radiative fluxes (e.g.,
 326 **Ramanathan et al. 1989; Ceppi et al. 2017**) – as a function of surface air temperature anomaly.
 327 The models disagree in the overall sign, as expected from CMIP5 models on shorter time scales
 328 (e.g., Vial et al. 2013; Caldwell et al. 2015), but can even change sign within a single simulation
 329 (e.g., ECEARTH or CESM *abrupt8x*). The strength of variation in time within one simulation
 330 can depend strongly on the forcing level (e.g. MIROC32 *1pct2x* vs. *1pct4x*) and the time scales
 331 of change differ between the models (e.g. IPSLCM5R vs. MPIESM12 *abrupt4x*). For some
 332 simulations, cloud response barely changes with temperature, contributing negligibly to the overall
 333 feedback (e.g. MPIESM12 *abrupt16x*, CESM104 *abrupt4x*, and MIROC32 *1pct2x*).

4. Discussion and Outlook

a. Published millennial-length simulations

Models of intermediate complexity are the most common tools used to study century to millennial time scales in the climate system (e.g., Zickfeld et al. 2013; Eby et al. 2013; Levermann et al. 2013; Rugenstein et al. 2016c; Jansen et al. 2018). However, they usually have a poorly resolved atmosphere and little or no representation of cloud processes. In contrast, the publications in Table 3 feature millennium-length AO-GCM simulations. Asterisks mark contributions to LongRunMIP. These papers provide a solid body of work on millennial-length climate simulations, but rarely use the same forcing levels and simulation length and focus on different aspects of the climate system. Three papers compare model formulation and processes of two AO-GCMs each (Frölicher et al. 2014; Paynter et al. 2018; Krasting et al. 2018), but otherwise models have not been systematically compared against each other. Fig. 4 and 7 show that AO-GCMs can strongly differ in their behavior. Spatial patterns of e.g., precipitation and surface heat fluxes also vary strongly between models and between different forcing scenarios for the same model (not shown), suggesting that some mechanisms and processes discussed in the published literature are not generalizable across models. For example, there is disagreement about which regions are thought to dominate the changing feedback parameter (Senior and Mitchell 2000; Andrews et al. 2015; Meraner et al. 2013; Caballero and Huber 2013) or whether or not, and on which time scales, the AMOC recovers from its initial reduction (Voss and Mikolajewicz 2001; Stouffer and Manabe 2003; Li et al. 2013; Rind et al. 2018; Thomas and Fedorov 2019). Paleo climate simulations are often several thousand years long, however, they usually include boundary conditions such as ice sheets or changing continental configurations, which differ from the ones used here. However, paleo climate studies often discuss equilibration time scales and deep ocean temperature trends relevant to the types

357 of models included in LongRunMIP (e.g., Brandefelt and Otto-Bliesner 2009; Zhang et al. 2013;
358 Klockmann et al. 2016; Marzocchi and Jansen 2017; Gottschalk et al. 2019).

359 *b. Limitations*

360 LongRunMIP analyses are currently limited mainly by the collected *variables* (Table 1). In-
361 cluding cloud fields and 3D atmospheric temperature and humidity fields, **for example**, would
362 allow users to study atmospheric dynamics and radiative feedbacks in more detail. The *differ-*
363 *ent forcing scenarios* of model contributions to LongRunMIP are both a strength and weakness.
364 Minimal requirements **have** encouraged a large number of contributions so far. However, study-
365 ing a single forcing scenario requires model selection or scaling between different forcing levels.
366 *Slab ocean simulations*, which replace a model’s dynamical ocean with a much shallower non-
367 dynamical mixed-layer, are a computationally cheap tool to compare fast and slow time scales and
368 the relevance of surface warming patterns (Boer and Yu 2003c; Danabasoglu and Gent 2009; Li
369 et al. 2013). We hope to receive submissions of these simulations in the future, to allow analysis of
370 their utility. Century to millennial-time scales in the real world include more processes and *Earth*
371 *System Feedbacks* than are included in LongRunMIP simulations, such as the carbon cycle, vege-
372 tation feedbacks, forcing agents other than CO₂ (such as other greenhouse gases or aerosols), ice
373 sheets, glacial rebound effects, changes to continental configuration, and orbital variation. Further,
374 the real climate system is never in equilibrium or steady state, because the forcing continuously
375 changes (e.g., Köhler et al. 2017). These Earth system feedbacks and additional forcings must be
376 taken into account when comparing the LongRunMIP models with paleo proxies or when project-
377 ing or predicting changes in future centuries or millennia.

378 *c. Summary and expected impact*

379 LongRunMIP is the first archive of millennial-length simulations of complex climate models,
380 featuring 50 simulations of 15 models by 10 modeling centers under various forcing scenarios (Ta-
381 ble 2). The archive provides an unprecedented opportunity to study the equilibrium response of a
382 large number of models to forcing. The variables included allow study of a range of phenomena
383 associated with the atmosphere, ocean, land, and sea ice (Table 1), and we expect LongRunMIP to
384 contribute to current discussions laid out in Section 1. This includes ocean heat uptake, sea level
385 rise, ocean circulation response to warming, large scale modes of variability, sea ice reduction,
386 polar amplification, precipitation variability, atmospheric dynamics, **long-term memory in time**
387 **series**, spatial warming patterns, ocean - atmosphere interactions, model spin-up techniques, the
388 relation of internal variability and forced response under different forcing levels, committed cli-
389 mate response, and the relation of time and state dependence of fast feedbacks and Earth System
390 Feedbacks and processes.

391 LongRunMIP is a MIP of opportunity, without an agreed upon protocol, and is a result of the
392 willingness of individual research groups to provide model output from simulations often con-
393 ducted over years of real-world time. As a result, the experiments are not standardized, but most
394 models provided a millennial-length simulation that begins with an abrupt quadrupling of CO₂
395 concentration. In addition to collecting simulations, we provide output with standardized formats
396 and variable names, and include versions regridded to a common grid, as well as global averages.

397 LongRunMIP builds upon a body of pioneering studies that looked at the behavior of models be-
398 yond the centennial scale (Table 3), LongRunMIP allows this sort of analysis to be applied across
399 a diverse group of models that exhibit strikingly different behavior (Fig. 7), and hopefully encour-

age others to look beyond the limitations and assumptions normally imposed by computational constraints, to directly study the equilibration of the fully coupled atmosphere-ocean system.

Data access and sharing

LongRunMIP currently consists of 15 TB of data and available for download at <https://data.iac.ethz.ch/longrunmip/>. Fields shown in this paper can be accessed on <https://data.iac.ethz.ch/longrunmip/BAMS/>.

See www.longrunmip.org for more details on available variables, contact information, sample figures and videos, and links to join a discussion community. We will be collecting more simulations over the next couple of years.

Acknowledgments. MR is funded by the Alexander von Humboldt Foundation. TA was supported by the Joint UK BEIS/Defra Met Office Hadley Centre Climate Programme (GA01101). NCAR is a major facility sponsored by the US National Science Foundation under Cooperative Agreement No. 1852977 TLF acknowledges support from the Swiss National Science Foundation under grant PP00P2_170687, from the EU-H2020 project CCiCC, and from the Swiss National Supercomputing Centre (CSCS). CL was supported through the Clusters of Excellence CliSAP (EXC177) and CLICCS (EXC2037), University Hamburg, funded through the German Research Foundation (DFG). SY was partly supported by European Research Council under the European Community's Seventh Framework Programme (FP7/20072013)/ERC grant agreement 610055 as part of the ice2ice project. This work was made possible for IPSL thanks to the HPC resources of TGCC and IDRIS made available by GENCI (Grand Equipement National de Calcul Intensif), CEA (Commissariat à l'Energie Atomique et aux Energies Alternatives) and CNRS (Centre National de la Recherche Scientifique) (project 016178).

References

- Andrews, T., J. M. Gregory, and M. J. Webb, 2015: The dependence of radiative forcing and feedback on evolving patterns of surface temperature change in climate models. *Journal of Climate*, **28** (4), 1630–1648, URL <http://dx.doi.org/10.1175/JCLI-D-14-00545.1>.
- Andrews, T., and M. J. Webb, 2018: The Dependence of Global Cloud and Lapse Rate Feedbacks on the Spatial Structure of Tropical Pacific Warming. *Journal of Climate*, **31** (2), 641–654, doi:10.1175/JCLI-D-17-0087.1, URL <https://doi.org/10.1175/JCLI-D-17-0087.1>.
- Andrews, T., and Coauthors, 2018: Accounting for Changing Temperature Patterns Increases Historical Estimates of Climate Sensitivity. *Geophysical Research Letters*, **45** (16), 8490–8499, doi:10.1029/2018GL078887, URL <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018GL078887>.
- Armour, K. C., 2017: Energy budget constraints on climate sensitivity in light of inconstant climate feedbacks. *Nature Climate Change*, **7**, 331 EP –, URL <http://dx.doi.org/10.1038/nclimate3278>.
- Armour, K. C., C. M. Bitz, and G. H. Roe, 2013: Time-Varying Climate Sensitivity from Regional Feedbacks. *Journal of Climate*, **26** (13), 4518–4534, URL <http://dx.doi.org/10.1175/JCLI-D-12-00544.1>.
- Armour, K. C., J. Marshall, J. R. Scott, A. Donohoe, and E. R. Newsom, 2016: Southern ocean warming delayed by circumpolar upwelling and equatorward transport. *Nature Geosci*, **9** (7), 549–554, URL <http://dx.doi.org/10.1038/ngeo2731>.

442 Bi, D., W. F. Budd, A. C. Hirst, and X. Wu, 2001: Collapse and reorganisation of the Southern
 443 Ocean overturning under global warming in a coupled model. *Geophysical Research Letters*,
 444 **28 (20)**, 3927–3930, URL <http://dx.doi.org/10.1029/2001GL013705>.

445 Bilbao, R. A., J. M. Gregory, and N. Bouttes, 2015: Analysis of the regional pattern of sea
 446 level change due to ocean dynamics and density change for 1993–2099 in observations and
 447 CMIP5 AOGCMs. *Climate Dynamics*, **45 (9-10)**, 2647–2666, URL <http://dx.doi.org/10.1007/s00382-015-2499-z>.

449 Bloch-Johnson, J., R. T. Pierrehumbert, and D. S. Abbot, 2015: Feedback temperature dependence
 450 determines the risk of high warming. *Geophysical Research Letters*, **42 (12)**, 4973– 4980, doi:
 451 10.1002/2015GL064240, URL <http://dx.doi.org/10.1002/2015GL064240>, 2015GL064240.

452 Bloch-Johnson, J., M. Rugenstein, and D. S. Abbot, in revision: Spatial radiative feedbacks from
 453 interannual variability using multiple regression. *Journal of Climate*.

454 Block, K., and T. Mauritsen, 2013: Forcing and feedback in the MPI-ESM-LR coupled model
 455 under abruptly quadrupled CO₂. *Journal of Advances in Modeling Earth Systems*, **5 (4)**, 676–
 456 691, URL <http://dx.doi.org/10.1002/jame.20041>.

457 Boer, G., and B. Yu, 2003a: Climate sensitivity and climate state. *Climate Dynamics*, **21 (2)**,
 458 167–176, URL <http://dx.doi.org/10.1007/s00382-003-0323-7>.

459 Boer, G., and B. Yu, 2003b: Climate sensitivity and response. *Climate Dynamics*, **20 (4)**, 415–429,
 460 URL <http://dx.doi.org/10.1007/s00382-002-0283-3>.

461 Boer, G. J., and B. Yu, 2003c: Dynamical aspects of climate sensitivity. *Geophysical Research*
 462 *Letters*, **30 (3)**, doi:10.1029/2002GL016549, URL <http://dx.doi.org/10.1029/2002GL016549>.

Brandefelt, J., and B. L. Otto-Bliesner, 2009: Equilibration and variability in a last
glacial maximum climate simulation with ccs3. *Geophysical Research Letters*, **36** (19),
doi:10.1029/2009GL040364, URL <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2009GL040364>.

Brown, P. T., Y. Ming, W. Li, and S. A. Hill, 2017: Change in the magnitude and mechanisms
of global temperature variability with warming. *Nature Climate Change*, **7**, 743 EP –, URL
<http://dx.doi.org/10.1038/nclimate3381>.

Burls, N. J., and A. V. Fedorov, 2017: Wetter subtropics in a warmer world: Contrasting past and
future hydrological cycles. *Proceedings of the National Academy of Sciences*, **114** (49), 12 888–
12 893, doi:10.1073/pnas.1703421114, URL <https://www.pnas.org/content/114/49/12888>.

Caballero, R., and M. Huber, 2013: State-dependent climate sensitivity in past warm climates and
its implications for future climate projections. *Proceedings of the National Academy of Sciences
of the United States of America*, **110** (35), 14 162–14 167, URL <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3761583/>.

Caldwell, P. M., M. D. Zelinka, K. E. Taylor, and K. Marvel, 2015: Quantifying the sources
of inter-model spread in equilibrium climate sensitivity. *Journal of Climate*, doi:10.1175/
JCLI-D-15-0352.1, URL <http://dx.doi.org/10.1175/JCLI-D-15-0352.1>.

Cao, L., L. Duan, G. Bala, and K. Caldeira, 2016: Simulated long-term climate response to
idealized solar geoengineering. *Geophysical Research Letters*, URL <http://dx.doi.org/10.1002/2016GL068079>.

Castruccio, S., D. J. McInerney, M. L. Stein, F. Liu Crouch, R. L. Jacob, and E. J. Moyer, 2014:
Statistical Emulation of Climate Model Projections Based on Precomputed GCM Runs. *Journal*

of *Climate*, **27** (5), 1829–1844, doi:10.1175/JCLI-D-13-00099.1, URL <https://doi.org/10.1175/JCLI-D-13-00099.1>, <https://doi.org/10.1175/JCLI-D-13-00099.1>.

Ceppi, P., F. Brient, M. D. Zelinka, and D. L. Hartmann, 2017: Cloud feedback mechanisms and their representation in global climate models. *Wiley Interdisciplinary Reviews: Climate Change*, **8** (4), e465, doi:10.1002/wcc.465, URL <https://onlinelibrary.wiley.com/doi/abs/10.1002/wcc.465>, <https://onlinelibrary.wiley.com/doi/pdf/10.1002/wcc.465>.

Ceppi, P., and J. M. Gregory, 2017: Relationship of tropospheric stability to climate sensitivity and earth’s observed radiation budget. *Proceedings of the National Academy of Sciences*, **114** (50), 13 126–13 131, doi:10.1073/pnas.1714308114, URL <https://www.pnas.org/content/114/50/13126>.

Charney, J., and Coauthors, 1979: Carbon Dioxide and Climate: A Scientific Assessment. Tech. rep., National Academy of Science, Washington, DC.

Clark, P. U., and Coauthors, 2016: Consequences of twenty-first-century policy for multi-millennial climate and sea-level change. *Nature Climate Change*, **6**, 360 EP –, URL <http://dx.doi.org/10.1038/nclimate2923>.

Collins, W. J., and Coauthors, 2011: Development and evaluation of an Earth-System model – HadGEM2. *Geoscientific Model Development*, **4** (4), 1051–1075, URL <https://www.geosci-model-dev.net/4/1051/2011/>.

Colman, R., and L. Hanson, 2016: On the relative strength of radiative feedbacks under climate variability and change. *Climate Dynamics*, 1–15, doi:10.1007/s00382-016-3441-8, URL <http://dx.doi.org/10.1007/s00382-016-3441-8>.

506 Colman, R., and B. McAvaney, 2009: Climate feedbacks under a very broad range of forcing.
 507 *Geophysical Research Letters*, **36** (1), n/a–n/a, doi:10.1029/2008GL036268, URL [http://dx.doi.](http://dx.doi.org/10.1029/2008GL036268)
 508 [org/10.1029/2008GL036268](http://dx.doi.org/10.1029/2008GL036268).

509 Cox, P. M., R. A. Betts, C. D. Jones, S. A. Spall, and I. J. Totterdell, 2000: Acceleration of global
 510 warming due to carbon-cycle feedbacks in a coupled climate model. *Nature*, **408**, 184 EP –,
 511 URL <http://dx.doi.org/10.1038/35041539>.

512 Danabasoglu, G., and P. R. Gent, 2009: Equilibrium Climate Sensitivity: Is It Accurate to Use
 513 a Slab Ocean Model? *Journal of Climate*, **22** (9), 2494–2499, URL [http://dx.doi.org/10.1175/](http://dx.doi.org/10.1175/2008JCLI2596.1)
 514 [2008JCLI2596.1](http://dx.doi.org/10.1175/2008JCLI2596.1).

515 Danabasoglu, G., S. G. Yeager, Y.-O. Kwon, J. J. Tribbia, A. S. Phillips, and J. W. Hurrell, 2012:
 516 Variability of the Atlantic Meridional Overturning Circulation in CCSM4. *Journal of Climate*,
 517 URL <http://dx.doi.org/10.1175/JCLI-D-11-00463.1>.

518 Deser, C., A. Phillips, V. Bourdette, and H. Teng, 2012: Uncertainty in climate change pro-
 519 jections: the role of internal variability. *Climate Dynamics*, **38** (3), 527–546, doi:10.1007/
 520 s00382-010-0977-x, URL <https://doi.org/10.1007/s00382-010-0977-x>.

521 Donner, L. J., and Coauthors, 2011: The Dynamical Core, Physical Parameterizations, and
 522 Basic Simulation Characteristics of the Atmospheric Component AM3 of the GFDL Global
 523 Coupled Model CM3. *Journal of Climate*, **24** (13), 3484–3519, URL [https://doi.org/10.1175/](https://doi.org/10.1175/2011JCLI3955.1)
 524 [2011JCLI3955.1](https://doi.org/10.1175/2011JCLI3955.1).

525 Dufresne, J.-L., and Coauthors, 2013: Climate change projections using the IPSL-CM5 Earth
 526 System Model: from CMIP3 to CMIP5. *Climate Dynamics*, **40** (9), 2123–2165, URL <https://doi.org/10.1007/s00382-012-1636-1>.
 527 [/doi.org/10.1007/s00382-012-1636-1](https://doi.org/10.1007/s00382-012-1636-1).

528 Dunne, J. P., and Coauthors, 2012: GFDL's ESM2 global coupled climate-carbon Earth System
 529 Models Part I: Physical formulation and baseline simulation characteristics. *Journal of Climate*.

530 Eby, M., and Coauthors, 2013: Historical and idealized climate model experiments: an inter-
 531 comparison of Earth system models of intermediate complexity. *Climate of the Past*, **9** (3),
 532 1111–1140, URL <http://www.clim-past.net/9/1111/2013/>.

533 Eyring, V., S. Bony, G. A. Meehl, C. A. Senior, B. Stevens, R. J. Stouffer, and K. E. Taylor,
 534 2016: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental
 535 design and organization. *Geoscientific Model Development*, **9** (5), 1937–1958, doi:10.5194/
 536 gmd-9-1937-2016, URL <https://www.geosci-model-dev.net/9/1937/2016/>.

537 Frölicher, T., and F. Joos, 2010: Reversible and irreversible impacts of greenhouse gas emissions in
 538 multi-century projections with the NCAR global coupled carbon cycle-climate model. *Climate*
 539 *Dynamics*, **35** (7-8), 1439–1459, doi:10.1007/s00382-009-0727-0, URL [http://dx.doi.org/10.](http://dx.doi.org/10.1007/s00382-009-0727-0)
 540 [1007/s00382-009-0727-0](http://dx.doi.org/10.1007/s00382-009-0727-0).

541 Frölicher, T. L., and D. J. Paynter, 2015: Extending the relationship between global warming and
 542 cumulative carbon emissions to multi-millennial timescales. *Environmental Research Letters*,
 543 **10** (7), 075 002, URL <http://stacks.iop.org/1748-9326/10/i=7/a=075002>.

544 Frölicher, T. L., M. Winton, and J. L. Sarmiento, 2014: Continued global warming after
 545 CO₂ emissions stoppage. *Nature Clim. Change*, **4** (1), 40–44, URL [http://dx.doi.org/10.1038/](http://dx.doi.org/10.1038/nclimate2060)
 546 [nclimate2060](http://dx.doi.org/10.1038/nclimate2060).

547 Gebbie, G., and P. Huybers, 2019: The Little Ice Age and 20th-century deep Pacific cooling.
 548 *Science*, **363** (6422), 70–74, doi:10.1126/science.aar8413, URL [http://science.sciencemag.org/](http://science.sciencemag.org/content/363/6422/70)
 549 [content/363/6422/70](http://science.sciencemag.org/content/363/6422/70).

550 Gent, P. R., and Coauthors, 2011: The Community Climate System Model Version 4. *Journal of*
551 *Climate*, **24** (19), 4973–4991, URL <http://dx.doi.org/10.1175/2011JCLI4083.1>.

552 Geoffroy, O., and D. Saint-Martin, 2014: Pattern decomposition of the transient climate response.
553 *Tellus A: Dynamic Meteorology and Oceanography*, **66** (1), 23 393, doi:10.3402/tellusa.v66.
554 23393, URL <https://doi.org/10.3402/tellusa.v66.23393>.

555 Geoffroy, O., D. Saint-Martin, G. Bellon, A. Voldoire, D. Olivié, and S. Tytéca, 2013a: Transient
556 Climate Response in a Two-Layer Energy-Balance Model. Part II: Representation of the Ef-
557 ficacy of Deep-Ocean Heat Uptake and Validation for CMIP5 AOGCMs. *Journal of Climate*,
558 **26** (6), 1859–1876, URL <http://dx.doi.org/10.1175/JCLI-D-12-00196.1>.

559 Geoffroy, O., D. Saint-Martin, D. J. L. Olivié, A. Voldoire, G. Bellon, and S. Tytéca, 2013b:
560 Transient Climate Response in a Two-Layer Energy-Balance Model. Part I: Analytical Solution
561 and Parameter Calibration Using CMIP5 AOGCM Experiments. *Journal of Climate*, **26** (6),
562 1841–1857, URL <http://dx.doi.org/10.1175/JCLI-D-12-00195.1>.

563 Gillett, N. P., V. K. Arora, K. Zickfeld, S. J. Marshall, and W. J. Merryfield, 2011: Ongoing
564 climate change following a complete cessation of carbon dioxide emissions. *Nature Geosci*,
565 **4** (2), 83–87, URL <http://dx.doi.org/10.1038/ngeo1047>.

566 Good, P., T. Andrews, R. Chadwick, J.-L. Dufresne, J. M. Gregory, J. A. Lowe, N. Schaller,
567 and H. Shiogama, 2016: nonlinMIP contribution to CMIP6: model intercomparison project
568 for non-linear mechanisms: physical basis, experimental design and analysis principles (v1.0).
569 *Geoscientific Model Development*, **9** (11), 4019–4028, doi:10.5194/gmd-9-4019-2016, URL
570 <https://www.geosci-model-dev.net/9/4019/2016/>.

571 Good, P., J. Gregory, J. Lowe, and T. Andrews, 2013: Abrupt CO₂ experiments as tools for pre-
 572 dicting and understanding CMIP5 representative concentration pathway projections. *Climate*
 573 *Dynamics*, **40** (3-4), 1041–1053, URL <http://dx.doi.org/10.1007/s00382-012-1410-4>.

574 Good, P., and Coauthors, 2015: Nonlinear regional warming with increasing co2 concentrations.
 575 *Nature Clim. Change*, **5** (2), 138–142, URL <http://dx.doi.org/10.1038/nclimate2498>.

576 Gottschalk, J., and Coauthors, 2019: Mechanisms of millennial-scale atmospheric co2 change
 577 in numerical model simulations. *Quaternary Science Reviews*, **220**, 30 – 74, doi:<https://doi.org/10.1016/j.quascirev.2019.05.013>, URL <http://www.sciencedirect.com/science/article/pii/S0277379118310473>.
 578
 579

580 Gregory, J. M., 2000: Vertical heat transports in the ocean and their effect on time-dependent cli-
 581 mate change. *Climate Dynamics*, **16**, 501–515, URL <http://dx.doi.org/10.1007/s003820000059>,
 582 [10.1007/s003820000059](http://dx.doi.org/10.1007/s003820000059).

583 Gregory, J. M., T. Andrews, and P. Good, 2015: The inconstancy of the transient climate re-
 584 sponse parameter under increasing CO₂. *Philosophical Transactions of the Royal Society of*
 585 *London A: Mathematical, Physical and Engineering Sciences*, **373** (2054), URL <http://rsta.royalsocietypublishing.org/content/373/2054/20140417>.
 586

587 Gregory, J. M., T. Andrews, P. Good, T. Mauritsen, and P. M. Forster, 2016: Small global-mean
 588 cooling due to volcanic radiative forcing. *Climate Dynamics*, 1–13, URL <http://dx.doi.org/10.1007/s00382-016-3055-1>.
 589

590 Gregory, J. M., and Coauthors, 2004: A new method for diagnosing radiative forcing and
 591 climate sensitivity. *Geophysical Research Letters*, **31** (3), URL <http://dx.doi.org/10.1029/2003GL018747>.
 592

Gregory, J. M., and Coauthors, 2005: A model intercomparison of changes in the Atlantic thermohaline circulation in response to increasing atmospheric CO₂ concentration. *Geophys. Res. Lett.*, **32** (L12703).

Hasumi, H., and S. Emori, 2004: K-1 Coupled GCM (MIROC) Description. Tech. rep., Center for Climate System Research CCSR, University of Tokyo, National Institute for Environmental Studies (NIES), Frontier Research Center for Global Change (FRCGC).

Hazeleger, W., and Coauthors, 2012: EC-Earth V2.2: description and validation of a new seamless earth system prediction model. *Climate Dynamics*, **39** (11), 2611–2629, URL <https://doi.org/10.1007/s00382-011-1228-5>.

Held, I., M. Winton, K. Takahashi, T. L. Delworth, F. Zeng, and G. Vallis, 2010: Probing the Fast and Slow Components of Global Warming by Returning Abruptly to Preindustrial Forcing. *Journal of Climate*, **23**, 2418 – 2427.

Hobbs, W., M. D. Palmer, and D. Monselesan, 2016: An Energy Conservation Analysis of Ocean Drift in the CMIP5 Global Coupled Models. *Journal of Climate*, **29** (5), 1639–1653, doi:10.1175/JCLI-D-15-0477.1, URL <http://dx.doi.org/10.1175/JCLI-D-15-0477.1>.

Jansen, M. F., L.-P. Nadeau, and T. M. Merlis, 2018: Transient versus Equilibrium Response of the Ocean’s Overturning Circulation to Warming. *Journal of Climate*, **31** (13), 5147–5163, doi:10.1175/JCLI-D-17-0797.1, URL <https://doi.org/10.1175/JCLI-D-17-0797.1>.

Jonko, A. K., K. M. Shell, B. M. Sanderson, and G. Danabasoglu, 2013: Climate Feedbacks in CCSM3 under Changing CO₂ Forcing. Part II: Variation of Climate Feedbacks and Sensitivity with Forcing. *Journal of Climate*, **26** (9), 2784–2795, URL <http://dx.doi.org/10.1175/JCLI-D-12-00479.1>.

615 Joshi, M., and J. Gregory, 2008: Dependence of the land-sea contrast in surface cli-
 616 mate response on the nature of the forcing. *Geophysical Research Letters*, **35** (24),
 617 doi:10.1029/2008GL036234, URL [https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2008GL036234)
 618 2008GL036234.

619 Jungclaus, J. H., and Coauthors, 2006: Ocean Circulation and Tropical Variability in the Coupled
 620 Model ECHAM5/MPI-OM. *Journal of Climate*, **19** (16), 3952–3972, URL [https://doi.org/10.](https://doi.org/10.1175/JCLI3827.1)
 621 1175/JCLI3827.1.

622 Khon, V. C., B. Schneider, M. Latif, W. Park, and C. Wengel, 2018: Evolution of Eastern Equa-
 623 torial Pacific Seasonal and Interannual Variability in Response to Orbital Forcing During the
 624 Holocene and Eemian From Model Simulations. *Geophysical Research Letters*, **45** (18), 9843–
 625 9851, doi:10.1029/2018GL079337, URL [https://agupubs.onlinelibrary.wiley.com/doi/abs/10.](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018GL079337)
 626 1029/2018GL079337.

627 Klockmann, M., U. Mikolajewicz, and J. Marotzke, 2016: The effect of greenhouse gas concentra-
 628 tions and ice sheets on the glacial AMOC in a coupled climate model. *Climate of the Past*, **12** (9),
 629 1829–1846, doi:10.5194/cp-12-1829-2016, URL <https://www.clim-past.net/12/1829/2016/>.

630 Knutti, R., 2010: The end of model democracy? *Climatic Change*, **102**, 395–404.

631 Knutti, R., M. A. A. Rugenstein, and G. C. Hegerl, 2017: Beyond equilibrium climate sensitivity.
 632 *Nature Geoscience*, **10**, 727 EP –, URL <http://dx.doi.org/10.1038/ngeo3017>.

633 Köhler, P., C. Nehrbass-Ahles, J. Schmitt, T. F. Stocker, and H. Fischer, 2017: A 156 kyr smoothed
 634 history of the atmospheric greenhouse gases CO₂, CH₄, and N₂O and their radiative forcing.
 635 *Earth System Science Data*, **9** (1), 363–387, doi:10.5194/essd-9-363-2017, URL [https://www.](https://www.earth-syst-sci-data.net/9/363/2017/)
 636 earth-syst-sci-data.net/9/363/2017/.

637 Kostov, Y., K. C. Armour, and J. Marshall, 2014: Impact of the atlantic meridional over-
638 turning circulation on ocean heat storage and transient climate change. *Geophysical Re-*
639 *search Letters*, **41** (6), 2108–2116, doi:10.1002/2013GL058998, URL [http://dx.doi.org/10.](http://dx.doi.org/10.1002/2013GL058998)
640 1002/2013GL058998.

641 Krasting, J. P., R. J. Stouffer, S. M. Griffies, R. W. Hallberg, S. L. Malyshev, B. L. Samuels,
642 and L. T. Sentman, 2018: Role of Ocean Model Formulation in Climate Response Uncertainty.
643 *Journal of Climate*, **31** (22), 9313–9333, doi:10.1175/JCLI-D-18-0035.1, URL [https://doi.org/](https://doi.org/10.1175/JCLI-D-18-0035.1)
644 10.1175/JCLI-D-18-0035.1, <https://doi.org/10.1175/JCLI-D-18-0035.1>.

645 Levermann, A., P. U. Clark, B. Marzeion, G. A. Milne, D. Pollard, V. Radic, and A. Robinson,
646 2013: The multimillennial sea-level commitment of global warming. *Proceedings of the Na-*
647 *tional Academy of Sciences*, **110** (34), 13 745–13 750, URL [http://www.pnas.org/content/110/](http://www.pnas.org/content/110/34/13745.abstract)
648 34/13745.abstract.

649 Li, C., J.-S. Storch, and J. Marotzke, 2013: Deep-ocean heat uptake and equilibrium cli-
650 mate response. *Climate Dynamics*, **40** (5-6), 1071–1086, URL [http://dx.doi.org/10.1007/](http://dx.doi.org/10.1007/s00382-012-1350-z)
651 s00382-012-1350-z.

652 Luo, Y., J. Lu, F. Liu, and O. Garuba, 2017: The Role of Ocean Dynamical Thermostat in De-
653 laying the El Niño–Like Response over the Equatorial Pacific to Climate Warming. *Journal*
654 *of Climate*, **30** (8), 2811–2827, doi:10.1175/JCLI-D-16-0454.1, URL [https://doi.org/10.1175/](https://doi.org/10.1175/JCLI-D-16-0454.1)
655 JCLI-D-16-0454.1.

656 Lutsko, N. J., and K. Takahashi, 2018: What Can the Internal Variability of CMIP5 Models Tell
657 Us about Their Climate Sensitivity? *Journal of Climate*, **31** (13), 5051–5069, doi:10.1175/
658 JCLI-D-17-0736.1, URL <https://doi.org/10.1175/JCLI-D-17-0736.1>.

- 659 Maher, N., D. Matei, S. Milinski, and J. Marotzke, 2018: Enso change in climate projec-
660 tions: Forced response or internal variability? *Geophysical Research Letters*, **45** (20), 11,390–
661 11,398, doi:10.1029/2018GL079764, URL [https://agupubs.onlinelibrary.wiley.com/doi/abs/10.](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018GL079764)
662 1029/2018GL079764.
- 663 Maher, N., and Coauthors, 2019: The Max Planck Institute Grand Ensemble: Enabling the
664 Exploration of Climate System Variability. *Journal of Advances in Modeling Earth Systems*,
665 **0** (0), doi:10.1029/2019MS001639, URL [https://agupubs.onlinelibrary.wiley.com/doi/abs/10.](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019MS001639)
666 1029/2019MS001639.
- 667 Manabe, S., R. J. Stouffer, M. J. Spelman, and K. Bryan, 1991: Transient Responses of a Cou-
668 pled Ocean Atmosphere Model to Gradual Changes of Atmospheric CO₂. Part I. Annual Mean
669 Response. *Journal of Climate*, **4** (8), 785–818.
- 670 Marzocchi, A., and M. F. Jansen, 2017: Connecting antarctic sea ice to deep-ocean circula-
671 tion in modern and glacial climate simulations. *Geophysical Research Letters*, **44** (12), 6286–
672 6295, doi:10.1002/2017GL073936, URL [https://agupubs.onlinelibrary.wiley.com/doi/abs/10.](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017GL073936)
673 1002/2017GL073936.
- 674 Mauritsen, T., and R. Pincus, 2017: Committed warming inferred from observations. *Nature Cli-*
675 *mate Change*, **7**, 652 EP –, URL <https://doi.org/10.1038/nclimate3357>.
- 676 Mauritsen, T., and Coauthors, 2018: Developments in the mpi-m earth system model version 1.2
677 (mpi-esm 1.2) and its response to increasing co2. *Journal of Advances in Modeling Earth Sys-*
678 *tems*, **0** (ja), doi:10.1029/2018MS001400, URL [https://agupubs.onlinelibrary.wiley.com/doi/](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018MS001400)
679 [abs/10.1029/2018MS001400](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018MS001400).

680 Meehl, G. A., C. Covey, T. Delworth, M. Latif, B. McAvaney, J. F. B. Mitchell, R. J. Stouffer,
681 and K. E. Taylor, 2007: THE WCRP CMIP3 Multimodel Dataset: A New Era in Climate
682 Change Research. *Bulletin of the American Meteorological Society*, **88** (9), 1383–1394, doi:
683 10.1175/BAMS-88-9-1383, URL <https://doi.org/10.1175/BAMS-88-9-1383>.

684 Meraner, K., T. Mauritsen, and A. Voigt, 2013: Robust increase in equilibrium climate sensitivity
685 under global warming. *Geophysical Research Letters*, **40** (22), 5944–5948, URL <http://dx.doi.org/10.1002/2013GL058118>.

686

687 Miller, R. L., and Coauthors, 2014: CMIP5 historical simulations (1850–2012) with GISS Mod-
688 elE2. *Journal of Advances in Modeling Earth Systems*, **6** (2), 441–478, URL <http://dx.doi.org/10.1002/2013MS000266>.

689

690 Nazarenko, L., and Coauthors, 2015: Future climate change under RCP emission scenarios with
691 GISS ModelE2. *Journal of Advances in Modeling Earth Systems*, **7** (1), 244–267, URL <http://dx.doi.org/10.1002/2014MS000403>.

692

693 Paynter, D., T. L. Frölicher, L. W. Horowitz, and L. G. Silvers, 2018: Equilibrium Climate
694 Sensitivity Obtained From Multimillennial Runs of Two GFDL Climate Models. *Journal of*
695 *Geophysical Research: Atmospheres*, **123** (4), 1921–1941, doi:10.1002/2017JD027885, URL
696 <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JD027885>.

697 Proistosescu, C., and P. J. Huybers, 2017: Slow climate mode reconciles historical and model-
698 based estimates of climate sensitivity. *Science Advances*, **3** (7), doi:10.1126/sciadv.1602821,
699 URL <http://advances.sciencemag.org/content/3/7/e1602821>.

700 Ramanathan, V., R. D. Cess, E. F. Harrison, P. Minnis, B. R. Barkstrom, E. Ahmad,
701 and D. Hartmann, 1989: Cloud-Radiative Forcing and Climate: Results from the Earth

702 Radiation Budget Experiment. *Science*, **243 (4887)**, 57–63, doi:10.1126/science.243.4887.
 703 57, URL <https://science.sciencemag.org/content/243/4887/57>, [https://science.sciencemag.org/](https://science.sciencemag.org/content/243/4887/57.full.pdf)
 704 [content/243/4887/57.full.pdf](https://science.sciencemag.org/content/243/4887/57.full.pdf).

705 Rehfeld, K., T. Münch, S. L. Ho, and T. Laepple, 2018: Global patterns of declining temperature
 706 variability from the Last Glacial Maximum to the Holocene. *Nature*, **554**, 356 EP –, URL <https://doi.org/10.1038/nature25454>.
 707

708 Rind, D., G. A. Schmidt, J. Jonas, R. Miller, L. Nazarenko, M. Kelley, and J. Romanski, 2018:
 709 Multicentury Instability of the Atlantic Meridional Circulation in Rapid Warming Simula-
 710 tions With GISS ModelE2. *Journal of Geophysical Research: Atmospheres*, **123 (12)**, 6331–
 711 6355, doi:10.1029/2017JD027149, URL [https://agupubs.onlinelibrary.wiley.com/doi/abs/10.](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2017JD027149)
 712 [1029/2017JD027149](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2017JD027149).

713 Rodgers, K. B., J. Lin, and T. L. Frölicher, 2015: Emergence of multiple ocean ecosystem drivers
 714 in a large ensemble suite with an Earth system model. *Biogeosciences*, **12 (11)**, 3301–3320,
 715 doi:10.5194/bg-12-3301-2015, URL <https://www.biogeosciences.net/12/3301/2015/>.

716 Rohrschneider, T., B. Stevens, and T. Mauritsen, 2019: On simple representations of the climate
 717 response to external radiative forcing. *Climate Dynamics*, doi:10.1007/s00382-019-04686-4.

718 Rugenstein, M., and Coauthors, 2019: Equilibrium climate sensitivity estimated by equilibrating
 719 climate models. *in revision for GRL*.

720 Rugenstein, M. A. A., K. Caldeira, and R. Knutti, 2016a: Dependence of global radiative feed-
 721 backs on evolving patterns of surface heat fluxes. *Geophysical Research Letters*, **43 (18)**, 9877–
 722 9885, URL <http://dx.doi.org/10.1002/2016GL070907>.

723 Rugenstein, M. A. A., J. M. Gregory, N. Schaller, J. Sedláček, and R. Knutti, 2016b: Multiannual
 724 Ocean–Atmosphere Adjustments to Radiative Forcing. *Journal of Climate*, **29** (15), 5643–5659,
 725 URL <http://dx.doi.org/10.1175/JCLI-D-16-0312.1>.

726 Rugenstein, M. A. A., J. Sedláček, and R. Knutti, 2016c: Nonlinearities in patterns of long-term
 727 ocean warming. *Geophysical Research Letters*, **43** (7), 3380–3388, URL [http://dx.doi.org/10.](http://dx.doi.org/10.1002/2016GL068041)
 728 [1002/2016GL068041](http://dx.doi.org/10.1002/2016GL068041).

729 Saint-Martin, D., and Coauthors, 2019: Fast forward to perturbed equilibrium climate.
 730 *Geophysical Research Letters*, **0** (ja), doi:10.1029/2019GL083031, URL [https://agupubs.](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019GL083031)
 731 [onlinelibrary.wiley.com/doi/abs/10.1029/2019GL083031](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019GL083031), [https://agupubs.onlinelibrary.wiley.](https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019GL083031)
 732 [com/doi/pdf/10.1029/2019GL083031](https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019GL083031).

733 Salzmann, M., 2017: The polar amplification asymmetry: role of Antarctic surface height.
 734 *Earth System Dynamics*, **8** (2), 323–336, doi:10.5194/esd-8-323-2017, URL [https://www.](https://www.earth-syst-dynam.net/8/323/2017/)
 735 [earth-syst-dynam.net/8/323/2017/](https://www.earth-syst-dynam.net/8/323/2017/).

736 Scheff, J., R. Seager, H. Liu, and S. Coats, 2017: Are Glacials Dry? Consequences for Pa-
 737 leoclimatology and for Greenhouse Warming. *Journal of Climate*, **30** (17), 6593–6609, doi:
 738 [10.1175/JCLI-D-16-0854.1](https://doi.org/10.1175/JCLI-D-16-0854.1), URL <https://doi.org/10.1175/JCLI-D-16-0854.1>.

739 Schmidt, G. A., and Coauthors, 2014: Configuration and assessment of the GISS ModelE2 contri-
 740 butions to the CMIP5 archive. *Journal of Advances in Modeling Earth Systems*, **6** (1), 141–184,
 741 URL <http://dx.doi.org/10.1002/2013MS000265>.

742 Schneider, T., C. M. Kaul, and K. G. Pressel, 2019: Possible climate transitions from breakup
 743 of stratocumulus decks under greenhouse warming. *Nature Geoscience*, **12** (3), 163–167, doi:
 744 [10.1038/s41561-019-0310-1](https://doi.org/10.1038/s41561-019-0310-1), URL <https://doi.org/10.1038/s41561-019-0310-1>.

- 745 Senior, C. A., and J. F. B. Mitchell, 2000: The time-dependence of climate sensitivity. *Geophysical*
746 *Research Letters*, **27** (17), 2685–2688, URL <http://dx.doi.org/10.1029/2000GL011373>.
- 747 Smith, R. S., J. M. Gregory, and A. Osprey, 2008: A description of the FAMOUS (version XD-
748 BUA) climate model and control run. *Geoscientific Model Development*, **1** (1), 53–68, doi:
749 10.5194/gmd-1-53-2008, URL <https://www.geosci-model-dev.net/1/53/2008/>.
- 750 Sniderman, J. M. K., and Coauthors, 2019: Southern Hemisphere subtropical drying as a transient
751 response to warming. *Nature Climate Change*, doi:10.1038/s41558-019-0397-9, URL <https://doi.org/10.1038/s41558-019-0397-9>.
752
- 753 Song, X., and G. J. Zhang, 2014: Role of Climate Feedback in El Nino-like SST Response to
754 Global Warming. *Journal of Climate*, doi:10.1175/JCLI-D-14-00072.1, URL [http://dx.doi.org/](http://dx.doi.org/10.1175/JCLI-D-14-00072.1)
755 10.1175/JCLI-D-14-00072.1.
- 756 Stouffer, R., and S. Manabe, 2003: Equilibrium response of thermohaline circulation to large
757 changes in atmospheric CO₂ concentration. *Climate Dynamics*, **20** (7-8), 759–773, URL <http://dx.doi.org/10.1007/s00382-002-0302-4>.
758
- 759 Stouffer, R. J., and S. Manabe, 1999: Response of a Coupled Ocean–Atmosphere Model to
760 Increasing Atmospheric Carbon Dioxide: Sensitivity to the Rate of Increase. *Journal of Cli-*
761 *mate*, **12** (8), 2224–2237, doi:10.1175/1520-0442(1999)012<2224:ROACOA>2.0.CO;2, URL
762 [http://dx.doi.org/10.1175/1520-0442\(1999\)012<2224:ROACOA>2.0.CO;2](http://dx.doi.org/10.1175/1520-0442(1999)012<2224:ROACOA>2.0.CO;2).
- 763 Svendsen, S. H., M. S. Madsen, Y. Suting, C. Rodehacke, and G. Adalgeirsdottir, 2015: An Intro-
764 duction to the Coupled EC-Earth-PISM Model System. report 15-05, Danish Climate Centre.

- 765 Taylor, K. E., R. J. Stouffer, and G. A. Meehl, 2011: An Overview of CMIP5 and the Ex-
766 periment Design. *Bulletin of the American Meteorological Society*, **93** (4), 485–498, doi:
767 10.1175/BAMS-D-11-00094.1, URL <http://dx.doi.org/10.1175/BAMS-D-11-00094.1>.
- 768 Thomas, M. D., and A. V. Fedorov, 2019: Mechanisms and Impacts of a Partial AMOC
769 Recovery Under Enhanced Freshwater Forcing. *Geophysical Research Letters*, **0** (0),
770 doi:10.1029/2018GL080442, URL [https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018GL080442)
771 2018GL080442.
- 772 Trossman, D. S., J. B. Palter, T. M. Merlis, Y. Huang, and Y. Xia, 2016: Large-scale ocean
773 circulation-cloud interactions reduce the pace of transient climate change. *Geophysical Re-*
774 *search Letters*, **43** (8), 3935–3943, URL <http://dx.doi.org/10.1002/2016GL067931>.
- 775 Vial, J., J.-L. Dufresne, and S. Bony, 2013: On the interpretation of inter-model spread in CMIP5
776 climate sensitivity estimates. *Climate Dynamics*, **41** (11-12), 3339–3362, URL [http://dx.doi.org/](http://dx.doi.org/10.1007/s00382-013-1725-9)
777 10.1007/s00382-013-1725-9.
- 778 Voldoire, A., and Coauthors, 2019: Evaluation of cmip6 deck experiments with cnrm-cm6-
779 1. *Journal of Advances in Modeling Earth Systems*, **0** (0), doi:10.1029/2019MS001683,
780 URL <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019MS001683>, <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019MS001683>.
- 782 Voss, R., and U. Mikolajewicz, 2001: Long-term climate changes due to increased CO2 con-
783 centration in the coupled atmosphere-ocean general circulation model ECHAM3/LSG. *Climate*
784 *Dynamics*, **17** (1), 45–60, doi:10.1007/PL00007925, URL <https://doi.org/10.1007/PL00007925>.
- 785 Winton, M., K. Takahashi, and I. M. Held, 2010: Importance of Ocean Heat Uptake Efficacy to
786 Transient Climate Change. *Journal of Climate*, **23** (9), 2333–2344, doi:10.1175/2009JCLI3139.

1, URL <http://dx.doi.org/10.1175/2009JCLI3139.1>.

Yamamoto, A., A. Abe-Ouchi, M. Shigemitsu, A. Oka, K. Takahashi, R. Ohgaito, and Y. Yamamaka, 2015: Global deep ocean oxygenation by enhanced ventilation in the Southern Ocean under long-term global warming. *Global Biogeochemical Cycles*, **29** (10), 1801–1815, URL <http://dx.doi.org/10.1002/2015GB005181>.

Yeager, S. G., C. A. Shields, W. G. Large, and J. J. Hack, 2006: The Low-Resolution CCSM3. *Journal of Climate*, **19** (11), 2545–2566, URL <http://dx.doi.org/10.1175/JCLI3744.1>.

Yoshimori, M., M. Watanabe, H. Shiogama, A. Oka, A. Abe-Ouchi, R. Ohgaito, and Y. Kamae, 2016: A review of progress towards understanding the transient global mean surface temperature response to radiative perturbation. *Progress in Earth and Planetary Science*, **3** (1), 21, doi:10.1186/s40645-016-0096-3, URL <https://doi.org/10.1186/s40645-016-0096-3>.

Zhang, X., G. Lohmann, G. Knorr, and X. Xu, 2013: Different ocean states and transient characteristics in last glacial maximum simulations and implications for deglaciation. *Climate of the Past*, **9** (5), 2319–2333, doi:10.5194/cp-9-2319-2013, URL <https://www.clim-past.net/9/2319/2013/>.

Zhou, C., M. D. Zelinka, and S. A. Klein, 2016: Impact of decadal cloud variations on the Earth’s energy budget. *Nature Geosci*, **9** (12), 871–874, URL <http://dx.doi.org/10.1038/ngeo2828>.

Zhu, J., Z. Liu, J. Zhang, and W. Liu, 2014: AMOC response to global warming: dependence on the background climate and response timescale. *Climate Dynamics*, **44** (11), 3449–3468, URL <http://dx.doi.org/10.1007/s00382-014-2165-x>.

Zickfeld, K., and Coauthors, 2013: Long-Term Climate Change Commitment and Reversibility: An EMIC Intercomparison. *Journal of Climate*, **26** (16), 5782–5809, URL <http://dx.doi.org/10.1175/JCLI-D-12-00584.1>.

809	LIST OF TABLES	
810	Table 1.	Description of collected variables. 2D means spatial resolution of latitude and
811		longitude, except for <i>msfmyz</i> where it means latitude and depth. 3D means lat-
812		itude, longitude, and depth. <i>msfmyz</i> is the sum of the eularian, eddybolus, and
813		submeso component. For <i>so</i> and <i>thetao</i> there are also February and September
814		values available for most models. 39
815	Table 2.	Overview of models and contributed simulations. The resolution of atmosphere
816		and ocean is given in # of grid points per latitude x longitude, and latitude x
817		longitude x depth, respectively. Models are referred to by their shortnames
818		throughout the manuscript. Section 2b describes the forcing levels. References
819		in the last column describe the models and simulations. Some simulations are
820		published in their full length, some simulations contributed to LongRunMIP are
821		the extensions of simulations discussed in the references, and some simulations
822		are unpublished. 40
823	Table 3.	Published millennial-length simulations 41

824 TABLE 1. Description of collected variables. 2D means spatial resolution of latitude and longitude, except for
825 *msftmyz* where it means latitude and depth. 3D means latitude, longitude, and depth. *msftmyz* is the sum of the
826 eularian, eddybolus, and submeso component. For *so* and *thetao* there are also February and September values
827 available for most models.

Shortname	Longname	Unit	Resolution
hfls	Surface Upward Latent Heat Flux	W m^{-2}	monthly, 2D
hfss	Surface Upward Sensible Heat Flux	W m^{-2}	monthly, 2D
pr	Precipitation on atmospheric grid	$\text{kg m}^{-2} \text{s}^{-1}$	monthly, 2D
psl	Sea Level Pressure	Pa	monthly, 2D
rlds	Surface Downwelling Longwave Radiation	W m^{-2}	monthly, 2D
rlus	Surface Upwelling Longwave Radiation	W m^{-2}	monthly, 2D
rlut	TOA Outgoing Longwave Radiation	W m^{-2}	monthly, 2D
rlutes	TOA Outgoing Clear-Sky Longwave Radiation	W m^{-2}	monthly, 2D
rsds	Surface Downwelling Shortwave Radiation	W m^{-2}	monthly, 2D
rsdt	TOA Incident Shortwave Radiation	W m^{-2}	monthly, 2D
rsus	Surface Upwelling Shortwave Radiation	W m^{-2}	monthly, 2D
rsut	TOA Outgoing Shortwave Radiation	W m^{-2}	monthly, 2D
rsutes	TOA Outgoing Clear-Sky Shortwave Radiation	W m^{-2}	monthly, 2D
tas	Near-Surface Air Temperature	K	monthly, 2D
ts	Atmospheric surface temperature	K	monthly, 2D
sic	Sea Ice Area Fraction	%	monthly, 2D
msftmyz	Meridional Overturning Circulation	$\text{m}^3 \text{s}^{-1}$	annual, 2D
tos	Sea surface temperature	K	annual, 2D
sos	Sea surface salinity	psu	annual, 2D
wfo	Net water flux into sea water	$\text{kg m}^{-2} \text{s}^{-1}$	annual, 2D
evs	Water evaporation	$\text{kg m}^{-2} \text{s}^{-1}$	annual, 2D
pr_ocn	Precipitation (rain and snow) on ocean grid	$\text{kg m}^{-2} \text{s}^{-1}$	annual, 2D
tauuo	Surface downward wind stress in x direction	N m^{-2}	annual, 2D
tauvo	Surface downward wind stress in y direction	N m^{-2}	annual, 2D
so	Sea Water Salinity	psu	annual, 3D
thetao	Sea Water Potential Temperature	K	annual, 3D

TABLE 2. Overview of models and contributed simulations. The resolution of atmosphere and ocean is given in # of grid points per latitude x longitude, and latitude x longitude x depth, respectively. Models are referred to by their shortnames throughout the manuscript. Section 2b describes the forcing levels. References in the last column describe the models and simulations. Some simulations are published in their full length, some simulations contributed to LongRunMIP are the extensions of simulations discussed in the references, and some simulations are unpublished.

Model (shortname)	Forcing level shortname	Length (yrs)	Atmosphere resolution	Ocean resolution	Control sim (yrs)	Model and simulation documentation
CCSM3 CCSM3	abrupt2x	3000	48 x 96	100 x 116 x 25	1530	Yeager et al. (2006) Danabasoglu and Gent (2009)
	abrupt4x	2120				
	abrupt8x	1450				
CCSM3 CCSM3II	abrupt2.4	3701	48 x 96	100 x 116 x 25	3805	Yeager et al. (2006) Castruccio et al. (2014)
	abrupt4.8	3132				
	lin2.4	3990				
CESM 1.0.4 CESM104	abrupt2x	2500	96 x 144	384 x 20 x 60	1320	Gent et al. (2011) Danabasoglu et al. (2012) Rugenstein et al. (2016c)
	abrupt4x	5900				
	abrupt8x	5100				
CNRM-CM6-1 CNRMCM61	abrupt2x	750	128 x 256	180 x 360 x 75	2000	Voldoire et al. (2019) Saint-Martin et al. (2019)
	abrupt4x	1850				
EC-Earth-PISM ECEARTH	historical RCP8.5+	1270	160 x 320	292 x 362 x 42	508	Hazeleger et al. (2012) Svendsen et al. (2015)
ECHAM5/MPIOM ECHAM5	abrupt4x	1000	48 x 96	101 x 120 x 40	100	Jungclaus et al. (2006) Li et al. (2013)
	1pct4x	6080				
FAMOUS FAMOUS	abrupt2x	3000	37 x 48	73 x 96 x 20	3000	Smith et al. (2008)
	abrupt4x	3000				
GFDL-CM3 GFDLCM3	1pct2x	5000	90 x 144	200 x 360 x 50	5200	Donner et al. (2011) Paynter et al. (2018)
GFDL-ESM2M GFDLESM2M	1pct2x	4500	90 x 144	200 x 360 x 50	1340	Dunne et al. (2012) Paynter et al. (2018)
GISS-E2-R GISSE2R	abrupt4x	5000	90 x 144	180 x 288 x 32	5225	Schmidt et al. (2014); Miller et al. (2014); Nazarenko et al. (2015) Rind et al. (2018)
	1pct4x	5000				
HadCM3L HadCM3L	abrupt2x	1000	73 x 96	73 x 96 x 20	1000	Cox et al. (2000) Cao et al. (2016)
	abrupt4x	1000				
	abrupt6x	1000				
	abrupt8x	1000				
HadGEM2-ES HadGEM2	abrupt4x	1328	145 x 192	216 x 360 x 40	239	Collins et al. (2011) Andrews et al. (2015)
IPSL-CM5A-LR IPSLCM5ALR	abrupt4x	1000	96 x 96	149 x 182 x 31	1000	Dufresne et al. (2013)
MIROC 3.2 MIROC32	1pct2x	2000	64 x 128	192 x 256 x 44	681	Hasumi and Emori (2004) Yamamoto et al. (2015); Yoshimori et al. (2016)
	1pct4x	2000				
MPIESM-1.2 MPIESM12	abrupt2x	1000	96 x 192	220 x 256 x 40	1237	Mauritsen et al. (2018) Rohrschneider et al. (2019)
	abrupt4x	1000				
	abrupt8x	1000				
	abrupt16x	1000				
MPIESM-1.1 MPIESM11	abrupt4x	4459	96 x 192	220 x 256 x 40	2000	Mauritsen et al. (2018)

TABLE 3. Published millennial-length simulations

Paper	Model	Forcing level	Length (yr)	Content/scientific comment
Senior and Mitchell (2000)	HadCM2	2xCO ₂	≈ 800	Included flux adjustments; effective climate sensitivity increases due to SW CRE due to changes in the inter-hemispheric temperature gradient
Bi et al. (2001)	CSIRO	3xCO ₂	≈ 1000	Cessation and recovery of Antarctic Bottom Water and North Atlantic Deep Water formation
Voss and Mikolajewicz (2001)	ECHAM3	2x and 4xCO ₂	850	Adjustment time scales, committed warming, ocean thermohaline circulation
Stouffer and Manabe (1999, GFDL 2003)	GFDL	0.5x, 2x, 4xCO ₂	and 4000	Thermohaline circulation and paleo-oceanographic implications
Boer and Yu (2003b,a,c)	CCCma	2050 and forcing	2100 1000	Radiative feedbacks and surface warming; effective climate sensitivity decreases with time; slab versus fully coupled models
Gregory et al. (2004)	HadCM3	2xCO ₂	≈ 1000	TOA radiative imbalance and surface temperature are not linearly related; after 1000 yr the model is still 0.7 W m ⁻² away from equilibrium
* Danabasoglu and Gent (2009)	CCSM3	2x, 4x, and 8xCO ₂	3000	Comparing slab and fully coupled models; determining ECS; Jonko et al. (2013) analyzed the contributions of different feedbacks to doublings of CO ₂
Gillett et al. (2011)	CanESM1	21st century	≈ 1000	Impact of reduced emissions
* Li et al. (2013)	ECHAM5/MPI-OM	2xCO ₂	≈ 6000	Comparing slab and fully coupled models; determining ECS; adjustment time scales of surface warming patterns, ocean heat uptake, and sea level rise
Frölicher et al. (2014); Frölicher and Paynter (2015)	GFDL-ESM2M, CSM1	4xCO ₂ pulse	1000	Climate impact of CO ₂ emission stoppage; evolving feedbacks; ECS; transient climate response to cumulative carbon emissions
* Andrews et al. (2015)	HadGEM2-ES	4xCO ₂	≈ 1300	Non-constancy of feedbacks; variations of TOA components cancel each other on the century to millennial time scale
* Yamamoto et al. (2015); Yoshimori et al. (2016)	MIROC3.2	2x and 4xCO ₂	2000	Deep ocean ventilation overall increases oxygenation after a transient decrease; review article on ocean heat uptake in coupled models and energy balance models
* Cao et al. (2016)	HadCM3L	2x, 4x, 6x, 8xCO ₂	1000	Comparing CO ₂ to other forcing agents and geo-engineering scenarios
* Rugenstein et al. (2016b,a)	CESM104	2x, 4x, 8xCO ₂	≈ 1300	Dependence of global and regional radiative feedback evolution on surface heat flux patterns; forcing adjustment
* Paynter et al. (2018)	GFDL-ESM2M, CM3	GFDL- 2xCO ₂	≈ 5000	Evolution of global and regional radiative feedbacks and the role of atmospheric vertical velocity fields and inversion strengths
* Rind et al. (2018)	GISS-E2-R	4xCO ₂	≈ 2000	AMOC reduction and recovery on North Atlantic surface flux conditions
Krasting et al. (2018)	GFDL-ESM2Mb, GFDL-ESM2G	4xCO ₂	5000	Ocean heat uptake, model formulation of diapycnal diffusivity and ocean vertical coordinates

LIST OF FIGURES

- Fig. 1.** Global **and annual** mean surface air temperature (*tas* in Table 1) anomaly and top of the atmosphere (TOA) radiative imbalance (computed as $rsdt - rlut - rsut$, see Table 1) to a step-forcing of quadrupling CO₂ as simulated by the CESM104 model. For the Coupled Model Intercomparison Project Phase 5 and 6, this simulation is part of the standard protocol, but only 150 simulated years are requested (blue shading). We collect simulations that extended this experiment for at least 850 years (light red shading), ideally until they are equilibrated (end of dark red shading). 43
- Fig. 2.** Global annual mean surface air temperature for all control (black) and forced (color, listed in the top right of each panel) simulations. *abrupt2x, 4x, 6x, 8x* means that the CO₂ concentration is doubled, quadrupled, sextupled, octupliated, as a step-forcing branched off the control simulation. *1pct2x* and *1pct4x* means the CO₂ concentration is linearly increased 1 % per year until the concentration is doubled or quadrupled, respectively. The simulations of ECEARTH and CCSM3II are described in Section b. Note the different axis ranges for each model. GFDLCM3 and CCSM3II are not branched off directly from the control simulation. 44
- Fig. 3.** Top of the atmosphere (TOA) annual and global mean radiative imbalance of all control simulations. Note the different lengths of the horizontal axes. The gray line indicates the average, the red line the linear trend, except for CCSM3II and GFDLCM3 for which both colors depict a fourth-order-polynomial fit. 45
- Fig. 4.** Global **and annual** mean temperature anomalies (experiment minus average of the control simulation) of the surface ocean (a, first layer) and deep ocean (b), as well as absolute values of deep ocean temperature in the control simulations (c), for *abrupt4x* (solid) and *1pct4x* (dashed) simulations. “Deep ocean” means around 2000 m **depth (closest level)**. Note that the time scale in c) is shorter than in a) and b). 46
- Fig. 5.** Time evolution of the surface air temperature anomaly in the *abrupt4x* simulations. The model mean of CCSM3, CESM104, CNRMCM61, ECHAM5, GISS2R, HadCM3L, HadGEM2, IPSLCM5A, MPIESM11, and MPIESM12 is shown in panel a, b, c, e, and f, while the model mean of only CESM104, GISS2R, and MPIESM11 is shown in panel d and g, due to the length of these contributions. See Table 2 for details of the length of each simulation. 47
- Fig. 6.** Time evolution of the zonal mean surface air temperature response normalized by the global mean temperature anomaly. Above (below) 1 means that warming is amplified (reduced) relative to the globally mean warming (a-d). Panel e-g show the differences (note the difference scale). Panel a, b, e, and f contain only *abrupt4x* simulations, while panel c, d, and g also contain the *1pct2x* and *RCP8.5+* simulations with integration lengths above 4000 years. Table 2 lists all simulations and model long names. 48
- Fig. 7.** Simulated shortwave cloud radiative effects **SW CRE** for different levels of global surface air temperature changes. Each point is a ten-year running average. Note the different axes labels, which cover a large range in TOA imbalance and surface temperature. Table 2 lists all simulations and model long names. 49






FIG. 1. Global **and annual** mean surface air temperature (*tas* in Table 1) anomaly and top of the atmosphere (TOA) radiative imbalance (computed as $rsdt - rlut - rsut$, see Table 1) to a step-forcing of quadrupling CO₂ as simulated by the CESM104 model. For the Coupled Model Intercomparison Project Phase 5 and 6, this simulation is part of the standard protocol, but only 150 simulated years are requested (blue shading). We collect simulations that extended this experiment for at least 850 years (light red shading), ideally until they are equilibrated (end of dark red shading).



881 FIG. 2. Global annual mean surface air temperature for all control (black) and forced (color, listed in the top
 882 right of each panel) simulations. *abrupt2x, 4x, 6x, 8x* means that the CO₂ concentration is doubled, quadrupled,
 883 sextupled, octupliated, as a step-forcing branched off the control simulation. *1pct2x* and *1pct4x* means the CO₂
 884 concentration is linearly increased 1 % per year until the concentration is doubled or quadrupled, respectively.
 885 The simulations of ECEARTH and CCSM3II are described in Section b. Note the different axis ranges for each
 886 model. GFDLCM3 and CCSM3II are not branched off directly from the control simulation.



887 FIG. 3. Top of the atmosphere (TOA) annual and global mean radiative imbalance of all control simulations.
 888 Note the different lengths of the horizontal axes. The gray line indicates the average, the red line the linear trend,
 889 except for CCSM3II and GFDLCM3 for which both colors depict a fourth-order-polynomial fit.






FIG. 4. Global and annual mean temperature anomalies (experiment minus average of the control simulation) of the surface ocean (a, first layer) and deep ocean (b), as well as absolute values of deep ocean temperature in the control simulations (c), for *abrup4x* (solid) and *1pct4x* (dashed) simulations. “Deep ocean” means around 2000 m depth (closest level). Note that the time scale in c) is shorter than in a) and b).


a) Temperature anomaly year 15-25





b) Temperature anomaly year 80-120




c) Temperature anomaly year 900-1000




d) Temperature anomaly year 4000-4200



e) Change in temperature anomalies (b)-(a)



f) Change in temperature anomalies (c)-(b)



g) Change in temperature anomalies (d)-(c)





FIG. 5. Time evolution of the surface air temperature anomaly in the *abrupt4x* simulations. The model mean of CCSM3, CESM104, CNRMCM61, ECHAM5, GISS2R, HadCM3L, HadGEM2, IPSLCM5A, MPIESM11, and MPIESM12 is shown in panel a, b, c, e, and f, while the model mean of only CESM104, GISS2R, and MPIESM11 is shown in panel d and g, due to the length of these contributions. See Table 2 for details of the length of each simulation.




FIG. 6. Time evolution of the zonal mean surface air temperature response normalized by the global mean temperature anomaly. Above (below) 1 means that warming is amplified (reduced) relative to the globally mean warming (a-d). Panel e-g show the differences (note the difference scale). Panel a, b, e, and f contain only *abrupt4x* simulations, while panel c, d, and g also contain the *1pct2x* and *RCP8.5+* simulations with integration lengths above 4000 years. Table 2 lists all simulations and model long names.




FIG. 7. Simulated shortwave cloud radiative effects **SW CRE** for different levels of global surface air temperature changes. Each point is a **ten-year** running average. **Note the different axes labels, which cover a large range in TOA imbalance and surface temperature.** Table 2 lists all simulations and model long names.